



Musculoskeletal Imaging Review Article

Tumors and pseudotumors of the soft tissues: Imaging semiology and strategy

Charlinne Paixao¹, Jean-Philippe Lustig¹, Sylvain Causeret², Loic Chaigneau³, Alexis Danner¹, Sebastien Aubry¹

¹Department Radiology, Musculoskeletal Imaging Unit, CHU Besancon, Besancon, ²Department of Surgery, CHU Dijon, Centre Georges-François Leclerc, Dijon, ³Department of Oncology, CHU Besancon, CHU de Besancon, Besancon, France.



***Corresponding author:**

Charlinne Paixao,
Department Radiology,
Musculoskeletal Imaging Unit,
CHU Besancon, Besancon,
France.

charlinne.paixao@orange.fr

Received : 07 August 2020
Accepted : 17 February 2021
Published : 03 March 2021

DOI
10.25259/JCIS_135_2020

Quick Response Code:



ABSTRACT

The aims of this educational review are to learn the semiological basis of soft-tissue lesions and, with the help of diagnostic algorithms, to apply the current recommendations for the management of soft-tissue tumors. Pseudotumors must first be identified and excluded. Among primary tumors, the search for macroscopic fat content on MRI is decisive; since it restricts the diagnostic range to adipocytic tumors. Key imaging features of non-adipocytic tumors are highlighted. When a deep soft-tissue mass is found, therapeutic abstention or simple monitoring is only appropriate when there is diagnostic certainty: This is only the case for typical pseudotumors, typical benign tumors, and fat tumors without atypical criteria. In all other cases, histological evidence is required. If there is any suspicion of soft-tissue sarcoma or any undetermined lesion, the patient should be referred to a sarcoma referral center before biopsy.

Keywords: Soft-tissue sarcoma, Adipocytic tumor, Non-adipocytic tumor, Lipoma, Liposarcoma

INTRODUCTION

The discovery of a soft-tissue mass is a common cause of requests for imaging. Ultrasound is the first-line imaging technique. However, an MRI is mandatory in the absence of a definite ultrasound diagnosis, or in case of a fascia or deep soft-tissue tumor. Malignant soft-tissue lesions, often soft-tissue sarcomas (STS), are rare but not exceptional (incidence \approx 3/100,000). Histological diagnosis is potentially difficult and often helped by cytogenetic data. The prognosis of malignant soft tissue tumors is all the more unfavorable when the diagnosis is delayed or when initial management is inadequate. The management of suspected soft-tissue tumors should always be discussed in a multidisciplinary meeting.

Pseudotumors must first be identified and eliminated. Among primary tumors, the search for a macroscopic fat component visible on MRI is decisive, since it restricts the diagnostic range to adipocytic tumors.

First, pseudotumors will be discussed. Then, most common characterizable tumors will be detailed by differentiating fatty from non-fatty tumors. Finally, a management strategy will be proposed using diagnostic algorithms [Figures 1-3] and based on current recommendations for the management of soft-tissue tumors.

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

©2020 Published by Scientific Scholar on behalf of Journal of Clinical Imaging Science




Figure 1: MRI diagnostic and management algorithm for lesions with spontaneous high signal on T1-weighted images.




Figure 2: MRI diagnostic algorithm for lesions with a liquid or pseudo-cystic high signal on T2-weighted images.




Figure 3: MRI diagnostic algorithm for low-signal lesions on T1-weighted and T2-weighted images.

PSEUDOTUMORS

Pseudotumors are lesions that mimic a tumor and that are not included in the WHO classification of soft-tissue tumors. Among the pseudotumors, intramuscular hematomas and soft-tissue abscesses both require caution and surveillance because of their sarcoma-like appearance.^[1,2]

Hematoma

The ultrasound similarity between a malignant tumor with necrosis and a hematoma should prompt the utmost caution [Figure 4a]. The presence of the following unfavorable factors should be investigated:^[3]

- Absence of trauma or mild trauma compared to the size of the hematoma
- Occurring in a child or teenager
- Intra-lesion vascular flow on color Doppler
- Absence of pain and slow progression, although these factors are less reliable because high-grade sarcomas are often painful and progress quickly.

If there is any doubt, an MRI scan [Figure 4b-d] and a biopsy should be performed. If a non-tumorous lesion is suspected, an imaging check-up at 3 weeks can be included in the clinical follow-up. Hematomas may show contrast enhancement, but only finely and peripherally [Figure 4d]. A nodular enhancement or an area of contrast enhancement in part of the lesion should suggest an underlying tumor and the biopsy should focus on that area.

Abscess

On ultrasound,^[4] an abscess appears as a heterogeneous collection, with hyperechoic peripheral infiltration related

to edema and peripheral hyperemia on color Doppler. If the abscess has a “falsely solid” appearance, mobilization shows fluctuation or displacement of its content. In the chronic stage, the liquefied center is anechogenic with posterior reinforcement [Figure 5a], the wall is echogenic and thick. Calcifications and air bubbles can be detected. Similarly, the ultrasound appearance of a high-grade sarcoma with necrosis may simulate an abscess. The absence of general clinical




Figure 4: A 32-year-old man with persistent pain in the left thigh 2 weeks after mild anterior trauma. US: Axial ultrasound image shows an anechoic collection with internal septa located in the vastus intermedius muscle (a). On MRI, the hematoma is well-defined, has a high-signal, and is surrounded by edema on fat-suppressed T2-weighted image (b), presents a low-signal deposit on gradient-echo T2-weighted image (c), and thin peripheral enhancement after injection of gadolinium chelates (d). A small post-traumatic lesion is also visible in the vastus lateralis muscle.




Figure 5: An 11-year-old girl with a popliteal fossa abscess consulting for pain and swelling with inflammatory syndrome. US: Axial ultrasound image of an abscess with hypoechoic and thick-walled, hyperechoic and heterogeneous content (a). MRI: On fat-saturated T1-weighted MRI image after injection of gadolinium chelates (b), the abscess, located between the semi-membranous and biceps muscles, has a low signal center, intense peripheral enhancement, and ill-defined peripheral enhancement due to perilesional edema.

and biological signs of infection should prompt caution. If there is any doubt, MRI and sampling of the lesion (possibly percutaneous microbiopsies under radiological control) should be performed.

On MRI, the center of an abscess typically has a low signal on T1-weighted (T1_w) images (although a high signal is possible), high signal on T2_w, and a high signal on diffusion-weighted with apparent diffusion coefficient (ADC) restriction.^[5] It has an outer shell that enhances after injection of contrast material^[6] and marked peri-lesional edema [Figure 5b]. Occasionally, a visible border in discrete high signal on T1_w images is indicative of granulation tissue.

Others pseudotumors

Others pseudotumors mainly include: Muscles and accessory muscle bundles, muscle hernia, Morel-Lavallee syndrome [Figure 6], circumscribed myositis ossificans [Figure 7], elastofibroma [Figure 8], synovial or mucoid cyst, cytosteatonecrosis, and parietal endometriosis following cesarean section and foreign body granuloma.

ADIPOCYTIC TUMORS

MRI is the reference method for the evaluation of deep fatty tumors. It is particularly discriminating when it enables the detection of macroscopic fat with high signal on T1_w and T2_w images, and low signal on sequences with fat signal suppression (Fat saturation, Water-only Dixon, Short Tau Inversion Recovery [STIR]) [Figure 1].

Lipoma

This is the most common soft-tissue tumor. Conventional lipoma is well defined, finely encapsulated, in a lobular arrangement. It is more often unilobular than liposarcoma.^[7] It presents a purely fatty signal in two thirds of cases [Figure 9a]. In one-third of cases, there is a nonfat component: calcifications, adipocyte necrosis, inflammatory infiltrate, vessels and phleboliths (angiolipoma), smooth muscle fibers (myolipoma), or myxoid areas.

Its septa are thin (<2 mm), and it does not show enhancement after contrast injection [Figure 9b]. Muscle fibers may be included in the lipoma, which is almost pathognomonic for benignity.

Liposarcoma

Liposarcoma is the second most common STS after undifferentiated pleomorphic cell sarcoma (14–18% of all soft-tissue malignancies). There are five major histologic types: Well-differentiated liposarcoma, dedifferentiated liposarcoma, myxoid liposarcoma, pleomorphic liposarcoma, and spindle cell liposarcoma.




Figure 6: A 22-year-old man with Morel-Lavallee syndrome who presented with left hip pain 4 weeks after a motorcycle accident. MRI: Collection with a low signal on T1-weighted (a) and high signal on T2-weighted (b) coronal images, between the hypodermic fat and the peripheral deep fascia of the left gluteus maximus muscle, typical of a Morel-Lavallee syndrome (arrows).




Figure 7: A 16-year-old girl presenting with increasing knee pain for 15 days and swelling. Natural evolution of a circumscribed myositis ossificans of the right thigh. MRI: Worrying pseudotumoral appearance on initial fat saturated proton density weighted axial MRI image (a). 7 months later on fat saturated T1-weighted axial MRI image after injection (b), the size of the lesion and peri-lesional infiltration decreased. The coronal CT reformat performed on the same day (c) shows ossification. Three years later, on fat saturated T1-weighted MRI image after injection (d), it has incorporated itself into the bone cortex.

Liposarcoma is extremely rare in neonates and children. About 85–90% of patients present a painless mass. Liposarcoma develops mainly at the expense of deep soft tissue, especially in the intermuscular spaces. Its two preferred sites are the thigh and the retroperitoneum. It is rare in the extremities (1–2%).




Figure 8: A 53-year-old woman with bilateral elastofibromas below the scapular tip, undergoing MRI for bilateral scapulargia. Typical laminated appearance on T1-weighted axial MRI image (arrows).




Figure 9: A 47-year-old woman with a conventional lipoma consulting for swelling of the right buttock. MRI: Lipoma of the right gluteus maximus muscle (arrow) showing a purely fatty signal on T1-weighted MRI image (a), low signal and no enhancement on fat saturated T1-weighted axial images after injection (b).

Calcifications are visible on radiography in 10% of cases.^[8] Ossifications are rare.

On CT and MRI, liposarcoma is a large tumor with polylobulated contours. It is quite limited, because it tends to grow by expansion rather than infiltration.^[7] Liposarcoma is usually multinodular in shape with small satellite nodules adjacent to the main tumor.^[9]

The appearance of liposarcoma depends on the degree of differentiation. The more differentiated the tumor, the fatter the signal: A well-differentiated liposarcoma can simulate a lipoma [Figure 10]. A signal intensity ratio on STIR images between the lesion and adjacent fat <1.4 and a maximum diameter <11 cm would be an additional argument in favor of a lipoma over atypical lipomatous tumors/well-differentiated liposarcomas.^[10] The slightest sign of atypia should lead to referral to a sarcoma center for image-guided percutaneous microbiopsy [Figure 1].^[11] A nodular non-fatty component of more than 2 cm in an atypical adipose tumor is in favor of a dedifferentiated liposarcoma.^[12] Pleomorphic, dedifferentiated, and myxoid liposarcomas may not contain a macroscopic fat component visible on imaging.^[13] Myxoid liposarcoma is




Figure 10: A 55-year-old man with liposarcoma consulting for swelling of the right thigh. MRI: T1-weighted (a) and fat saturated T2-weighted coronal MRI images (b). Fatty tumor of the right thigh whose signal is heterogeneous (arrows) and higher than that of subcutaneous fat (arrowhead). Biopsy confirmed the diagnosis of well-differentiated liposarcoma.

characterized by low density pseudocystic areas on CT scan, with a T1_w and T2_w pseudocystic signal on MRI that is intensely enhanced after contrast injection.^[14] Fatty areas account for less than 10% of the mass. If MRI is performed without contrast injection, myxoid liposarcoma may be confused with cystic lesions and operated inappropriately.

Hibernoma

Hibernoma is a very rare benign tumor that occurs in adults aged 20–40 years. It is composed of brown fat present in newborns that is similar to that of hibernating animals. Hibernoma is most frequently reported in the thigh, in the superficial soft tissues more than in deep tissue.^[15] Hibernoma is frequently discovered incidentally on PET-CT due to its high hypermetabolism.

On ultrasound, hibernoma is hyperechoic, but heterogeneous with hyperemia and arterial-like flows on Doppler.^[16]

On CT, its density is intermediate, between that of fat and muscle, and heterogeneous due to the presence of fibrous septa.

On MRI, it presents a fat signal different from that of subcutaneous fat: An intermediate signal between muscle and fat on T1_w images (“dirty white appearance”), slightly heterogeneous, close to that of fat on T2_w and intense in STIR.^[16] Fat saturation only partially suppresses its signal.^[17] The lesion is partitioned by typically central fibrous septa. The presence of large intra-tumoral vessels is an inconsistent but useful sign, since it is absent in liposarcomas^[18] [Figure 11]. There is a rare myxoid variant with a pseudo-liquid signal on T2_w images.

As the lesion is hypervascular, percutaneous biopsy presents a risk of hemorrhage.

Intramuscular hemangioma

Intramuscular hemangioma is a benign vascular tumor composed of capillary proliferation and vessels. This term should not be used for purely cavernous forms, which are vascular malformations. Intramuscular hemangioma often has a macroscopic fat component.^[19] Other signs are:

- On standard radiography, the presence of phleboliths
- On ultrasonography, a heterogeneous multi-cystic lesion with serpiginous images with very low or even undetectable flow on color Doppler
- On MRI, a polylobulated, well-delimited, heterogeneous lesion with predominantly high signal on T1_w and T2_w images, and the presence of intralesional vessels that are highly enhanced after injection.

NON-ADIPOCYTIC TUMORS

When the lesion does not contain a macroscopic fatty component, imaging can still help in the diagnosis, by showing a melanocytic [Figure 1], myxoid [Figure 2], or fibrous component [Figure 3].

Melaninic tumor: Clear cell sarcomas

This is a melanocytic sarcoma generally affecting women between the ages of 25 and 40. The preferred topography is the limbs (especially the lower limbs, at the foot), opposite a tendon, ligament, or fascia. It has the MRI signal of melanin in about 50% of cases [Figure 1]. It is often non-encapsulated, with blurred borders, homogeneous, without necrosis or peri-lesional edema, and enhances after injection.^[20]

Myxoid tumors

Myxoid tumors show pseudo-liquid high signal on T2_w images and enhancement after gadolinium injection [Figure 2]. In the presence of a deep soft-tissue mass presenting a myxoid signal, the following entities should be considered: Intramuscular myxoma [Figure 12], myxoid chondrosarcoma, myxofibrosarcoma [Figure 13], myxoid liposarcoma (see above), or intramuscular neurogenic tumor (schwannoma or neurofibroma).

Neurogenic tumors should be considered in the presence of a mass centered on a nerve that shows the split fat sign (thin layer of periwound fat) and the target sign (central fibrocellular zone and peripheral myxoid zone).^[21]

Fibrous tumors

Nodular fasciitis

Nodular fasciitis is the most common benign fibrous soft-tissue tumor. Its spontaneous growth is limited. It occurs




Figure 11: A 36-year-old woman with a hibernoma who presented with a subcutaneous mass of the left thigh. MRI appearance of a hibernoma of the posterior compartment of the left thigh on T1-weighted axial image (a), T2-weighted sagittal image (b), and fat saturated T1-weighted axial image after injection (c). The brown fat of the hibernoma (arrows) has a lower signal on T1-weighted and T2-weighted images than that of the subcutaneous fat. There are also large intralesional vessels enhanced after injection (arrowhead).



Figure 12: A 66-year-old woman with intramuscular myxoma who presented with palpable swelling of the right deltoid muscle. MRI: High T2-weighted signal and septa (a). It presents weak internal enhancement, with enhancement of its pseudo-capsule on fat saturated T1-weighted axial images after injection (b).




Figure 13: A 42-year-old man with myxofibrosarcoma consulting for a painful mass in the right thigh. MRI: On fat saturated T2-weighted axial image (a), this fibrosarcoma is very heterogeneous due to fibrous, myxoid, and tissue content, and areas of hemorrhagic necrosis. Therefore, on fat saturated T1-weighted axial image after injection (b), enhancement is also heterogeneous.

mostly in young adults and with no predominance in either sex. Clinically, it is a single mass of less than 3 cm, firm and painful, growing rapidly in 2–3 months, and it may disappear spontaneously. The nodule is always unique and mainly affects

the upper limb.^[22] Its location with regard to the fascia makes it possible to distinguish between subcutaneous, intramuscular (mimicking a sarcoma), or intermuscular forms.

On ultrasonography, nodular fasciitis is a hypoechoic mass that is often heterogeneous. On MRI, it is a homogeneous T1_w isosignal mass with a heterogeneous, and moderately high signal on T2_w images, which enhances diffusely and homogeneously or peripherally after gadolinium injection.^[23] The subcutaneous form is the most frequent, forming rounded nodules developed from the deep peripheral fascia and extending subcutaneously. There are often characteristic extensions that produce a “fascial tail sign,” an evocative but non-specific sign. Intermuscularly, the tumor develops in the fascia, often with star-shaped contours, mimicking an inflammatory, or aggressive lesion.^[22,24] It may have a central cystic zone, which is a sign of the “inverted target.”^[23,24]

Desmoid tumor

The desmoid tumor is part of the group of deep aggressive fibromatosis. Its location is deep, subaponeurotic, unlike superficial plantar and palmar fibromatoses. It affects all ages but especially young women. It presents slow and painless growth, which explains its large size (>10 cm). It is rarely multifocal and is limited to a single limb most of the time. It does not metastasize, but is locally aggressive.

It can be distinguished as follows:

- The abdominal parietal form, the most frequent, mainly affecting young women (context: Pregnancy or surgical scar)
- The extra-abdominal form occurring mainly at the root of the limbs, especially the shoulder^[25]
- And the intra-abdominal form.

The ultrasound appearance, which is not very specific, shows a poorly delimited hypoechoic mass with posterior acoustic shadowing and possible hypervascularization.

On MRI, the extra-abdominal form is typically intermuscular and can be surrounded by fat (“split fat sign”), like a neurogenic tumor. However, it also frequently infiltrates the muscles. Its signal is variable and often heterogeneous. Depending on the degree of maturation, it is hyperintense (young lesion with significant cellularity) or hypointense on T2_w images [chronic fibrous form rich in collagen, Figure 3].^[26,27] The presence of hypointense bands within the mass on all sequences is quite frequent and evocative (corresponding to dense collagen patches). The enhancement is moderate to high. Erosion or periosteal apposition of the facing bone is possible, but the medullary canal is not invaded. A linear extension of these lesions is often observed along the fascia, giving the appearance of a “deer antler” or “fascial tail sign” [Figure 14].^[28]

Because recurrences are common even after complete surgical removal, treatment options include monitoring, discontinuation of oral contraceptives, and possibly systemic therapies or cryoablation under imaging control.^[29]

Fibrosarcoma

Fibrosarcoma is a rare sarcoma that affects middle-aged subjects, and usually develops in the deep soft parts of the extremities, trunk or cervicofacial area. Its signal depends on its content (myxoid [Figure 2], fibrous [Figure 3], and necrotic).

Inflammatory myofibroblastic tumors

These tumors were previously referred to as “inflammatory pseudotumors.” Also known as plasma cell granulomas, they are rare neoplasms. They occur in children or young adults and can occur in any part of the body. They consist of a single or multinodular mass of variable size that may include tissular, myxoid, hemorrhagic, necrotic, or calcified areas. The treatment is surgical excision. Imaging is non-specific.

DISEASE MANAGEMENT

When a deep soft-tissue mass is found, therapeutic abstention or simple monitoring is only appropriate when

there is diagnostic certainty: This is only the case for typical pseudotumors, typical benign tumors, and fat tumors without atypical criteria measuring less than 10 cm [Figure 1]. In all other cases, imaging can suggest diagnostic hypotheses, but histological evidence by percutaneous microbiopsy under imaging control will be essential.

The essential elements of the imaging report

The report must include the following elements:

- Position of the lesion in relation to the deep peripheral fascia (i.e., superficial or deep)
- The exact compartmental location
- Relationships (vascular, nervous, bone, and joint)
- Size in all three dimensions
- Form and boundaries
- And the composition or matrix.

The conclusion must mention one or more diagnostic hypotheses, and specify if necessary the elements of subsequent management, for example, complementary further imaging examinations, monitoring, and/or discussion in a multidisciplinary meeting.

Ultrasound

As a first-line examination, ultrasound confirms the deep site of the lesion and provides semiological information.

Typical benign lesions that do not require further exploration are:^[30]

- Simple cysts
- Well-defined synovial bursae or cysts, with purely fluid content, without a solid component, anechogenic with posterior enhancement and without internal vascularization
- Superficial, homogeneous, well-defined, encapsulated, compressible, and stable for at least 6 months
- Vascular malformations, stable for at least 6 months
- Granulomas consistent with clinical history
- Muscular hernias
- And Morton’s neuromas.




Figure 14: A 30-year-old woman with a desmoid tumor who presented palpable swelling of the right leg. MRI: T1-weighted axial image (a), T2-weighted axial image (b), and fat saturated T1-weighted image after injection (c) of a desmoid tumor of the superficial posterior compartment of the right leg. It presents an intermediate signal with a central fibrous area (arrow) with low signal regardless of weighting. Although non-specific, its extension along the fascia (arrowheads), clearly visible after injection (c), is suggestive of the diagnosis.

Post-traumatic muscle injury requires caution and control at 3–6 weeks is essential.

Ultrasound “red flags” are as follows:

- Deep localization. A peripheral lesion in intimate contact with, or crossing the deep peripheral fascia is also suspect
- Anarchic vascularization
- Heterogeneity
- Areas of necrosis
- Hypoechogenicity.

If there is any doubt about the diagnosis, the ultrasound must be supplemented by MRI.^[21]

MRI

MRI indications are broad. MRI should be carried out immediately if there is a clinical suspicion of malignancy: Large, fixed or deep lesion, persistent post-traumatic swelling, increase in lesion size, or bone or joint damage. It is indicated as soon as a “red flag” is observed on ultrasound, when the size of the mass exceeds 5 cm, and for any recurrent tumor at an operated site.^[30]

MRI is the reference examination for the characterization of the tumor matrix according to its signal on T1_w [Figure 1], T2_w [Figure 2], T2*_w [Figure 3] images, and its enhancement after injection of contrast medium. The search for macroscopic fat component visible on MRI is decisive, since it restricts the diagnostic range to adipocytic tumors and a few benign tumors containing fat.

MRI is essential for the assessment of loco regional extension. The field of view must encompass the entire lesion and peripheral edema with a margin. A skin marker is useful. The analysis is essentially based on high-resolution axial sections. However, at least two orthogonal acquisition planes or 3D-acquisition are required to accurately measure the lesion.

All non-fatty tumors, purely fat tumors larger than 10 cm, and fat tumors with any atypical criteria require pathological evidence (except for genetic exceptions).^[11] In the case of a deep, purely fatty tumor of less than 10 cm, with a location that is “easy” for the surgeon to access, surveillance or surgery must be discussed in a multidisciplinary meeting. In addition to the criteria for atypical fat tumors [Figure 1], some MRI signs are suggestive of a malignant lesion.^[31,32]

- Heterogeneity on T2_w images (necrosis)
- Arterial enhancement after gadolinium injection^[32]
- Retroperitoneal location
- Erosion of the adjacent bone
- Spread to a vascular pedicle in contact.

The regularity or otherwise of the lesion limits is not a discriminating criterion.^[33]

Many studies have focused on diffusion imaging. The difference between an abscess [Figure 15] and a necrotic tumor [Figure 16] could be due to diffusion-weighted imaging. Both would have a high diffusion-weighted signal,




Figure 15: A 38-year-old man with osteitis and abscess who presented for left forearm pain without fever. T1-weighted MR image (a), fat-suppressed T2-weighted image (b), fat-suppressed T1-weighted image after injection of gadolinium chelates (c) and diffusion weighted image (d) of radial osteitis: Cortical erosion of low T1-weighted signal and high diffusion-weighted signal of the medullary. Deep muscular abscess (short radial extensor of the carpus, short extensor, and long abductor of the thumb) have low T1-weighted signal (a), high T2-weighted signal (b), peripheral enhancement (c), and high diffusion-weighted signal (d). Bacteriological analysis found *Staphylococcus aureus*.




Figure 16: A 72-year-old man with dedifferentiated spindle cell sarcoma who presented for growing abdominal wall mass. Fat-suppressed T2-weighted MR image (a), fat-suppressed T1-weighted image after gadolinium chelate injection (b), diffusion-weighted image (c) and Apparent Diffusion Coefficient (d). Moderately high diffusion-weighted image with high ADC of the necrotic portion and low ADC of the tissue portion.

but the abscess would have a low ADC, unlike the necrotic sarcoma, which would have a high ADC.

Beyond its usefulness in abscess, diffusion imaging would help distinguish benign from malignant tumors. The ADC of malignant tumors is lower than that of benign tumors. However, there is a large overlap, which precludes sufficient diagnostic reliability.^[34]

Multidisciplinary management and sampling recommendations

If there is any suspicion of STS and any undetermined lesion, the patient should be referred to a sarcoma referral center before biopsy.^[35] Percutaneous biopsy should be performed after MRI and the multidisciplinary meeting.^[36] The biopsy path, defined in agreement with the surgeon, must take into account the compartmental anatomy of the limb, so as not to transform an intracompartmental lesion into an extracompartmental lesion. It should be in the axis of the limbs or ribs. Numerous samples must be taken with a coaxial needle (≥ 16 Gauge) from the various lesion components, avoiding necrotic areas, and major vascular-nerve axes. A tattoo is sometimes performed to mark the biopsy site. In addition to pathological analysis, molecular biology analysis should also be systematically performed^[37] and centralized review in a sarcoma referral center.^[38] During the operation, the surgeon will have to resect the path and the mass in a single block to achieve complete macroscopic and microscopic margin-negative “R0” resection.

CONCLUSION

The discovery of a deep soft-tissue mass is a problem commonly encountered by radiologists. In line with the adage whereby “the ignorant asserts, the learned doubts, the wise thinks,” should be referred to a referral center for multidisciplinary care, in which oncologists, radiologists, pathologists, surgeons, and radiotherapists specialized in the management of STS will collaborate.

Declaration of patient consent

Patient’s consent not required as patients identity is not disclosed or compromised.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Laredo JD, Wybier M, Petrover D. *Imagerie Rhumatologique et Orthopédique*. Netherlands: Elsevier; 2016.
- Cotten A. *Pseudo-Tumeurs des Tissus Mous, in Imagerie Musculosquelettique: Pathologies Générales*. Netherlands: Elsevier, Masson; 2013.
- Taïeb S, Penel N, Vanseymortier L, Ceugnart L. Soft tissue sarcomas or intramuscular haematomas? *Eur J Radiol* 2009;72:44-9.
- Chau CL, Griffith JF. Musculoskeletal infections: Ultrasound appearances. *Clin Radiol* 2005;60:149-59.
- Costa FM, Ferreira EC, Vianna EM. Diffusion-weighted magnetic resonance imaging for the evaluation of musculoskeletal tumors. *Magn Reson Imaging Clin N Am* 2011;19:159-80.
- Turecki MB, Taljanovic MS, Stubbs AY, Graham AR, Holden DA, Hunter TB, *et al.* Imaging of musculoskeletal soft tissue infections. *Skeletal Radiol* 2010;39:957-71.
- Matsumoto K, Hukuda S, Ishizawa M, Chano T, Okabe H. MRI findings in intramuscular lipomas. *Skeletal Radiol* 1999;28:145-52.
- Murphey MD, Arcara LK, Fanburg-Smith J. From the archives of the AFIP: Imaging of musculoskeletal liposarcoma with radiologic-pathologic correlation. *Radiographics* 2005;25:1371-95.
- Jelinek JS, Kransdorf MJ, Shmookler BM, Abouafia AJ, Malawer MM. Liposarcoma of the extremities: MR and CT findings in the histologic subtypes. *Radiology* 1993;186:455-9.
- Donners R, Krieg AH, Baumhoer D, Boll DT, Harder D. Quantitative signal intensity ratios to distinguish between subfascial lipoma and atypical lipomatous tumor/well-differentiated liposarcoma using short-tau inversion recovery (STIR) MRI. *Diagn Interv Imaging* 2020;101:383-90.
- Fouque O, Kind M, Boulet B, Brisse H, Kemel S, Genah I, *et al.* Strategy for the diagnosis of an adipocytic soft tissue tumor of the adult. *J D'imagerie Diagnostique Interv* 2018;1:265-83.
- Murphey MD. World Health Organization classification of bone and soft tissue tumors: Modifications and implications for radiologists. *Semin Musculoskelet Radiol* 2007;11:201-14.
- Kransdorf MJ, Bancroft LW, Peterson JJ, Murphey MD, Foster WC, Temple HT. Imaging of fatty tumors: Distinction of lipoma and well-differentiated liposarcoma. *Radiology* 2002;224:99-104.
- Sung MS, Kang HS, Suh JS, Lee JH, Park JM, Kim JY, *et al.* Myxoid liposarcoma: Appearance at MR imaging with histologic correlation. *Radiographics* 2000;20:1007-19.
- Furlong MA, Fanburg-Smith JC, Miettinen M. The morphologic spectrum of hibernoma: A clinicopathologic study of 170 cases. *Am J Surg Pathol* 2001;25:809-14.
- Anderson SE, Schwab C, Stauffer E, Banic A, Steinbach LS. Hibernoma: Imaging characteristics of a rare benign soft tissue tumor. *Skeletal Radiol* 2001;30:590-5.
- Papathanassiou ZG, Alberghini M, Taïeb S, Errani C, Picci P, Vanel D. Imaging of hibernomas: A retrospective study on twelve cases. *Clin Sarcoma Res* 2011;1:1-12.
- Colville J, Feigin K, Antonescu CR, Panicek DM. Hibernoma: Report emphasizing large intratumoral vessels and high T1

- signal. *Skeletal Radiol* 2006;35:547-50.
19. Dubois J, Alison M. Vascular anomalies: What a radiologist needs to know. *Pediatr Radiol* 2010;40:895-905.
 20. de Beuckeleer LH, de Schepper AM, Vandevenne JE, Bloem JL, Davies AM, Oudkerk M, *et al.* MR imaging of clear cell sarcoma (malignant melanoma of the soft parts): A multicenter correlative MRI-pathology study of 21 cases and literature review. *Skeletal Radiol* 2000;29:187-95.
 21. Kransdorf MJ, Murphey MD. Imaging of soft-tissue musculoskeletal masses: Fundamental concepts. *Radiographics* 2016;36:1931-48.
 22. Polat P, Kantarci M, Alper F, Gursan N, Suma S, Okur A. Nodular fasciitis of the breast and knee in the same patient. *AJR Am J Roentgenol* 2002;178:1426-8.
 23. Coyle J, White LM, Dickson B, Ferguson P, Wunder J, Naraghi A. MRI characteristics of nodular fasciitis of the musculoskeletal system. *Skeletal Radiol* 2013;42:975-82.
 24. Wang XL, de Schepper AM, Vanhoenacker F, de Raeye H, Gielen J, Aparisi F, *et al.* Nodular fasciitis: Correlation of MRI findings and histopathology. *Skeletal Radiol* 2002;31:155-61.
 25. Pignatti G, Barbanti-Brodano G, Ferrari D, Gherlinzoni F, Bertoni F, Bacchini P, *et al.* Extraabdominal desmoid tumor: A study of 83 Cases. *Clin Orthop Relat Res* 2000;375:207-13.
 26. Casillas J, Sais GJ, Greve JL, Iparraguirre MC, Morillo G. Imaging of intra-and extraabdominal desmoid tumors. *Radiographics* 1991;11:959-68.
 27. Vandevenne JE, de Schepper AM, de Beuckeleer L, van Marck E, Aparisi F, Bloem JL, *et al.* New concepts in understanding evolution of desmoid tumors: MR imaging of 30 lesions. *Eur Radiol* 1997;7:1013-9.
 28. Robbin MR, Murphey MD, Temple HT, Kransdorf MJ, Choi JJ. Imaging of musculoskeletal fibromatosis. *Radiographics* 2001;21:585-600.
 29. Schmitz JJ, Schmit GD, Atwell TD, Callstrom MR, Kurup AN, Weisbrod AJ, *et al.* Percutaneous cryoablation of extraabdominal desmoid tumors: A 10-year experience. *AJR Am J Roentgenol* 2016;207:190-5.
 30. Noebauer-Huhmann IM, Weber MA, Lalam RK, Trattng S, Bohndorf K, Vanhoenacker F, *et al.* Soft tissue tumors in adults: ESSR-approved guidelines for diagnostic imaging. *Semin Musculoskelet Radiol* 2015;19:475-82.
 31. de Schepper AM, Ramon FA, Degryse HR. Statistical analysis of MRI parameters predicting malignancy in 141 soft tissue masses. *Röfo* 1992;156:587-91.
 32. Benedikt RA, Jelinek JS, Kransdorf MJ, Moser RP, Berrey BH. MR imaging of soft-tissue masses: Role of gadopentetate dimeglumine. *J Magn Reson Imaging* 1994;4:485-90.
 33. Peabody TD, Simon MA. Principles of staging of soft-tissue sarcomas. *Clin Orthop Relat Res* 1993;289:19-31.
 34. Dallaudière B, Lecouvet F, Berg BV, Omoumi P, Perlepe V, Cerny M, *et al.* Diffusion-weighted MR imaging in musculoskeletal diseases: Current concepts. *Diagn Interv Imaging* 2015;96:327-40.
 35. ESMO/European Sarcoma Network Working Group. Soft tissue and visceral sarcomas: ESMO clinical practice guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2014;25 Suppl 3:iii102-12.
 36. Heudel PE, Cousin P, Lurkin A, Cropet C, Ducimetiere F, Collard O, *et al.* Territorial inequalities in management and conformity to clinical guidelines for sarcoma patients: An exhaustive population-based cohort analysis in the Rhône-Alpes region. *Int J Clin Oncol* 2014;19:744-52.
 37. Ducimetiere F, Lurkin A, Ranchère-Vince D, Decouvelaere AV, Péoc'h M, Istier L, *et al.* Incidence of sarcoma histotypes and molecular subtypes in a prospective epidemiological study with central pathology review and molecular testing. *PLoS One* 2011;6:e20294.
 38. Ray-Coquard I, Montesco MC, Coindre JM, dei Tos AP, Lurkin A, Ranchère-Vince D, *et al.* Sarcoma: Concordance between initial diagnosis and centralized expert review in a population-based study within three European regions. *Ann Oncol* 2012;23:2442-9.

How to cite this article: Paixao C, Lustig J, Causeret S, Chaigneau L, Danner A, Aubry S. Tumors and pseudotumors of the soft tissues: Imaging semiology and strategy. *J Clin Imaging Sci* 2021;11:13.